

# Numerical Solution of Fifth Order Boundary Value Problems by Petrov-Galerkin Method with Quintic B-splines as basis functions and Septic B-Splines as weight functions

S. M. Reddy

Sreenidhi institute of science and technology, Department of Science and Humanities  
Yamnapet, Ghatkesar, Hyderabad  
Telangana, India-501 301  
smrmitw@gmail.com

**Abstract:** This paper deals with a finite element method involving Petrov-Galerkin method with quintic B-splines as basis functions and septic B-splines as weight functions to solve a general fifth order boundary value problem with a particular case of boundary conditions. The basis functions are redefined into a new set of basis functions which vanish on the boundary where the Dirichlet type of boundary conditions are prescribed. The weight functions are also redefined into a new set of weight functions which in number match with the number of redefined basis functions. The proposed method was applied to solve several examples of fifth order linear and nonlinear boundary value problems. The obtained numerical results were found to be in good agreement with the exact solutions available in the literature.

**Keywords:** Petrov-Galerkin method, Quintic B-spline, Septic B-spline, Fifth order boundary value problem, Absolute error.

## 1. Introduction

In this paper, we consider a general fourth order boundary value problem

$$a_0(x)y^{(5)}(x) + a_1(x)y^{(4)}(x) + a_2(x)y'''(x) + a_3(x)y''(x) + a_4(x)y'(x) + a_5(x)y(x) = b(x), c < x < d \quad (1)$$

subject to boundary conditions

$$y(c) = A_0, y(d) = C_0, y'(c) = A_1, y'(d) = C_1, y''(c) = A_2 \quad (2)$$

where  $A_0, A_1, A_2, C_0, C_1$  are finite real constants and

$$a_0(x), a_1(x), a_2(x), a_3(x), a_4(x), a_5(x) \text{ and } b(x)$$

are all continuous functions defined on the interval  $[c, d]$ .

The fifth order boundary value problems occur in the mathematical modelling of the viscoelastic flows and other branches of mathematical, physical and engineering sciences [1, 2]. The existence and uniqueness of the solution for these types of problems have been discussed in Agarwal [3]. Finding the analytical solutions of such type of boundary value problems in general is not possible. Over the years, many researchers have worked on boundary value problems by using different methods for numerical solutions [4-13]. Wazwaz [4] developed the solution of special type of fifth order boundary value problems by using the modified Adomain decomposition method and he provided the solution in the form of a rapidly convergent series, Siddiqi et al. [5] presented the solution of a special case of linear fifth order boundary value problems by using quartic spline functions, Rashidinia et al. [6] presented the solution of a special case of linear fifth order boundary value problem by using non-polynomial spline functions techniques, Noor and Sayed [7] applied the Homotopy perturbation method for solving fifth order boundary value problems, Caglar and Caglar [8] presented the Local polynomial regression method to solve the special case of fifth

order boundary value problems, Gamel [9] presented the solution of fifth order boundary value problems by Sinc-Galerkin method, Syam and Ahili [10] presented combination of Adomain decomposition method and the Homotopy method to solve a fifth order singularly perturbed boundary value problem arising in viscoelastic flows, Zhao [11] developed the solution of fifth order boundary value problems by variational iteration method, Lamnii et al. [12] developed the sextic spline collocation method to solve special case of fifth order boundary value problems, Kasi Viswanadham and Sreenivasulu [13] developed the quartic B-spline Galerkin method to a general solve fifth order boundary value problem. So far, fifth order boundary value problems have not been solved by using Petrov-Galerkin method with quintic B-splines as basis functions and septic B-splines as weight functions. This motivated us to solve a fifth order boundary value problem by Petrov-Galerkin method with quintic B-splines as basis functions and septic B-splines as weight functions.

In this paper, we try to present a simple finite element method which involves Petrov-Galerkin approach with quintic B-splines as basis functions and septic B-splines as weight functions to solve the fifth order boundary value problem of the type (1)-(2). This paper is organized as follows. Section 2, deals with the justification for using Petrov-Galerkin Method. In section 3, the definition of quintic B-splines and septic B-splines has been described. In section 4, description of the Petrov-Galerkin method with quintic B-splines as basis functions and septic B-splines as weight functions has been presented and in section 5, solution procedure to find the nodal parameters is presented. In section 6, the proposed method is tested on several linear and nonlinear boundary value problems. The solution to a nonlinear problem has been obtained as the limit of a sequence of solution of linear problems generated by the quasilinearization technique [14]. Finally, in the last section, the conclusions are presented.

## 2. Justification for using Petrov-Galerkin method

In Finite Element Method (FEM) the approximate solution can be written as a linear combination of basis functions which constitute a basis for the approximation space under consideration. FEM involves variational methods like Rayleigh Ritz method, Galerkin method, Least Squares method, Petrov-Galerkin method and Collocation method etc.

In Petrov-Galerkin method, the residual of approximation is made orthogonal to the weight functions. When we use Petrov-Galerkin method, a weak form of approximation solution for a given differential equation exists and is unique under appropriate conditions [15, 16] irrespective of properties of a given differential operator. Further, a weak solution also tends to a classical solution of given differential equation, provided sufficient attention is given to the boundary conditions [17]. That means the basis functions should vanish on the boundary where the Dirichlet type of boundary conditions are prescribed and also the number of weight functions should match with the number of basis functions. Hence in this paper we employed the use of Petrov-Galerkin method with quintic B-splines as basis functions and septic B-splines as weight functions to approximate the solution of fifth order boundary value problem.

## 3. Definition of quintic B-spline and Septic B-spline

The quintic B-splines and septic B-splines are defined in [18]-[20]. The existence of quintic spline interpolate  $s(x)$  to a function in a closed interval  $[c, d]$  for spaced knots (need not be evenly spaced) of a partition  $c = x_0 < x_1 < \dots < x_{n-1} < x_n = d$  is established by constructing it. The construction of  $s(x)$  is done with the help of the quintic B-splines. Introduce ten additional knots  $x_{-5}, x_{-4}, x_{-3}, x_{-2}, x_{-1}, x_{n+1}, x_{n+2}, x_{n+3}, x_{n+4}$  and  $x_{n+5}$  in such a way that

$$x_{-5} < x_{-4} < x_{-3} < x_{-2} < x_{-1} < x_0 \text{ and } x_n < x_{n+1} < x_{n+2} < x_{n+3} < x_{n+4} < x_{n+5}.$$

Now the quintic B-splines  $B_i(x)$ 's are defined by

$$B_i(x) = \begin{cases} \sum_{r=i-3}^{i+3} \frac{(x_r - x)_+^5}{\pi'(x_r)}, & x \in [x_{i-3}, x_{i+3}] \\ 0, & \text{otherwise} \end{cases}$$

where

$$(x_r - x)_+^5 = \begin{cases} (x_r - x)^5, & \text{if } x_r \geq x \\ 0, & \text{if } x_r \leq x \end{cases}$$

$$\text{and } \pi(x) = \prod_{r=i-3}^{i+3} (x - x_r)$$

where  $\{B_{-2}(x), B_{-1}(x), B_0(x), B_1(x), \dots, B_n(x), B_{n+1}(x), B_{n+2}(x)\}$  forms a basis for the space  $S_5(\pi)$  of quintic polynomial splines. Schoenberg [20] has proved that quintic B-splines are the unique nonzero splines of smallest compact support with the knots at

$$x_{-5} < x_{-4} < x_{-3} < x_{-2} < x_{-1} < x_0 < x_1 < \dots < x_{n-1} < x_n < x_{n+1} < x_{n+2} < x_{n+3} < x_{n+4} < x_{n+5}.$$

In a similar analogue septic B-splines  $R_i(x)$ 's are defined by

$$R_i(x) = \begin{cases} \sum_{r=i-4}^{i+4} \frac{(x_r - x)_+^7}{\pi'(x_r)}, & x \in [x_{i-4}, x_{i+4}] \\ 0, & \text{otherwise} \end{cases}$$

where

$$\text{and } \pi(x) = \prod_{r=i-4}^{i+4} (x - x_r)$$

where  $\{R_{-3}(x), R_{-2}(x), R_{-1}(x), R_0(x), R_1(x), \dots, R_{n-1}(x), R_n(x), R_{n+1}(x), R_{n+2}(x), R_{n+3}(x)\}$  forms a basis for the space  $S_7(\pi)$  of septic polynomial splines with the introduction of four more additional knots  $x_{-7}, x_{-6}, x_{n+6}, x_{n+7}$  to the already existing knots

$x_{-5}$  to  $x_{n+5}$ . Schoenberg [20] has proved that septic B-splines are the unique nonzero splines of smallest compact support with the knots at

$$x_{-7} < x_{-6} < x_{-5} < x_{-4} < x_{-3} < x_{-2} < x_{-1} < x_0 < x_1 < \dots$$

$$< x_{n-1} < x_n < x_{n+1} < x_{n+2} < x_{n+3} < x_{n+4} < x_{n+5} < x_{n+6} < x_{n+7}.$$

## 4. Description of the method

To solve the boundary value problem (1) subject to boundary conditions (2) by the Petrov-Galerkin method with quintic B-splines as basis functions and septic B-splines as weight functions, we define the approximation for  $y(x)$  as

$$y(x) = \sum_{j=-2}^{n+2} \alpha_j B_j(x) \quad (3)$$

where  $\alpha_j$ 's are the nodal parameters to be determined and

$B_j(x)$ 's are quintic B-spline basis functions. In Petrov-Galerkin method, the basis functions should vanish on the boundary where the Dirichlet type of boundary conditions are specified. In the set of quintic B-splines  $\{B_{-2}(x), B_{-1}(x), B_0(x), \dots, B_n(x), B_{n+1}(x), B_{n+2}(x)\}$ , the basis functions  $B_{-2}(x), B_{-1}(x), B_0(x), B_1(x), B_2(x), B_{n-2}(x), B_{n-1}(x), B_n(x), B_{n+1}(x)$  and  $B_{n+2}(x)$  do not vanish at one of the boundary points. So, there is a necessity of redefining the basis functions into a new set of basis functions which vanish on the boundary where the Dirichlet type of boundary conditions are specified. The procedure for redefining of the basis functions is as follows.

Using the definition of quintic B-splines and the Dirichlet boundary conditions of (2), we get the approximate solution at the boundary points as

$$y(c) = y(x_0) = \sum_{j=-2}^2 \beta_j B_j(x_0) = A_0 \quad (4)$$

$$y(d) = y(x_n) = \sum_{j=n-2}^{n+2} \beta_j B_j(x_n) = C_0 \quad (5)$$

Eliminating  $\alpha_{-2}$  and  $\alpha_{n+2}$  from the equations (3), (4) and (5), we get

$$y(x) = w(x) + \sum_{j=-1}^{n+1} \alpha_j P_j(x) \quad (6)$$

where

$$w(x) = \frac{A_0}{B_{-2}(x_0)} B_{-2}(x) + \frac{C_0}{B_{n+2}(x_n)} B_{n+2}(x) \quad (7)$$

$$P_j(x) = \begin{cases} B_j(x) - \frac{B_j(x_0)}{B_{-2}(x_0)} B_{-2}(x), & j = -1, 0, 1, 2 \\ B_j(x), & j = 3, 4, \dots, n-3 \\ B_j(x) - \frac{B_j(x_n)}{B_{n+2}(x_n)} B_{n+2}(x), & j = n-2, n-1, n, n+1 \end{cases} \quad (8)$$

The new set of basis functions in the approximation  $y(x)$  is  $\{P_j(x), j = -1, 0, \dots, n+1\}$ . Here  $w(x)$  takes care of given set of Dirichlet boundary conditions and  $P_j(x)$ 's vanish on the boundary. In Petrov-Galerkin method, the number of basis functions in the approximation should match with the number of weight functions. Here the number of basis functions in the approximation is  $n + 3$ , where as the number of weight functions is  $n+7$ . So, there is a need to redefine the weight functions into a new set of weight functions which in number match with the number of basis functions. The procedure for redefining the weight functions is as follows:

Let us write the approximation for  $v(x)$  as

$$v(x) = \sum_{j=-3}^{n+3} \beta_j R_j(x) \quad (9)$$

where  $R_j(x)$ 's are septic B-splines and here we assume that above approximation  $v(x)$  satisfies corresponding homogeneous boundary conditions of the given boundary conditions (2). That means  $v(x)$  defined in (9) satisfies the conditions

$$v(c)=0, v(d)=0, v'(c)=0, v'(d)=0 \quad (10)$$

Applying the boundary conditions (10) to (9), we get the approximate solution at the boundary points as

$$v(c) = v(x_0) = \sum_{j=-3}^3 \beta_j R_j(x_0) = 0 \quad (11)$$

$$v(d) = v(x_n) = \sum_{j=n-3}^{n+3} \beta_j R_j(x_n) = 0 \quad (12)$$

$$v'(c) = v'(x_0) = \sum_{j=-3}^{n+3} \beta_j R'_j(x_0) = 0 \quad (13)$$

$$v'(d) = v'(x_n) = \sum_{j=n-3}^{n+3} \beta_j R'_j(x_n) = 0 \quad (14)$$

Eliminating  $\beta_{-3}, \beta_{-2}, \beta_{n+2}$  and  $\beta_{n+3}$  from the equations (9) and (11) to (14), we get the approximation for  $v(x)$  as

$$v(x) = \sum_{j=-1}^{n+1} \beta_j T_j(x) \quad (15)$$

where

$$T_j(x) = \begin{cases} S_j(x) - \frac{S'_j(x_0)}{S'_{-2}(x_0)} S_{-2}(x), & j = -1, 0, 1, 2, 3 \\ S_j(x), & j = 4, 5, \dots, n-4 \\ S_j(x) - \frac{S'_j(x_n)}{S'_{n+2}(x_n)} S_{n+2}(x), & j = n-3, n-2, n-1, n, n+1 \end{cases} \quad (16)$$

$$S_j(x) = \begin{cases} R_j(x) - \frac{R_j(x_0)}{R_{-3}(x_0)} R_{-3}(x), & j = -2, -1, 0, 1, 2, 3 \\ R_j(x), & j = 4, 5, \dots, n-4 \\ R_j(x) - \frac{R_j(x_n)}{R_{n+3}(x_n)} R_{n+3}(x), & j = n-3, n-2, n-1, n, n+1, n+2 \end{cases} \quad (17)$$

Now the new set of weight functions for the approximation  $v(x)$  is  $\{T_j(x), j = -1, 0, \dots, n+1\}$ . Here  $T_j(x)$ 's and their derivatives vanish on the boundary.

Applying the Petrov-Galerkin method to (1) with the new set of basis functions  $\{P_j(x), j = -1, 0, \dots, n+1\}$  and with the new set of weight functions  $\{T_j(x), j = -1, 0, \dots, n+1\}$ , we get

$$\int_{x_0}^{x_n} [a_0(x)y^{(5)}(x) + a_1(x)y^{(4)}(x) + a_2(x)y'''(x) + a_3(x)y''(x) + a_4(x)y'(x) + a_5(x)y(x)] T_i(x) dx = \int_{x_0}^{x_n} b(x) T_i(x) dx \quad \text{for } i = -1, 0, \dots, n+1. \quad (18)$$

Integrating by parts the first term on the left hand side of (18) and after applying the boundary conditions prescribed in (2), we get

$$\begin{aligned} \int_{x_0}^{x_n} a_0(x)y^{(5)}(x) T_i(x) dx &= \frac{d^2}{dx^2} [a_0(x) T_i(x)]_{x_n} y''(x_n) \\ &- \frac{d^2}{dx^2} [a_0(x) T_i(x)]_{x_0} A_2 - \frac{d^3}{dx^3} [a_0(x) T_i(x)]_{x_n} C_1 \\ &+ \frac{d^3}{dx^3} [a_0(x) T_i(x)]_{x_0} A_1 + \int_{x_0}^{x_n} \frac{d^4}{dx^4} [a_0(x) T_i(x)] y'(x) dx \end{aligned} \quad (19)$$

Substituting (19) in (18) and using the approximation for  $y(x)$  given in (6), and after rearranging the terms for resulting equations, we get system of equations in the matrix form as

$$\mathbf{A} \boldsymbol{\alpha} = \mathbf{B} \quad (20)$$

where  $\mathbf{A} = [a_{ij}]$ ;

$$\begin{aligned} a_{ij} &= \int_{x_0}^{x_n} \{a_1(x) T_i(x) P_j^{(4)}(x) + a_2(x) T_i(x) P_j'''(x) + a_3(x) T_i(x) P_j''(x) \\ &+ [\frac{d^4}{dx^4} [a_0(x) T_i(x)] + a_4(x) T_i(x)] P_j'(x) + a_5(x) T_i(x) P_j(x)\} dx \\ &+ \frac{d^2}{dx^2} [a_0(x) T_i(x)]_{x_n} P_j''(x_n) \end{aligned} \quad \text{for } i = -1, 0, \dots, n+1; j = -1, 0, \dots, n+1. \quad (21)$$

$\mathbf{B} = [b_i]$ ;

$$\begin{aligned}
 b_i = & \int_{x_0}^{x_n} \{b(x)T_i(x) - [a_1(x)T_i(x)w^{(4)}(x) + a_2(x)T_i(x)w'''(x) \\
 & + a_3(x)T_i(x)w''(x) + [\frac{d^4}{dx^4}[a_0(x)T_i(x)] + a_4(x)T_i(x)]w'(x) \\
 & + a_5(x)T_i(x)w(x)]\} dx - \frac{d^2}{dx^2}[a_0(x)T_i(x)]_{x_n} w''(x_n) \\
 & + \frac{d^2}{dx^2}[a_0(x)T_i(x)]_{x_0} A_2 + \frac{d^3}{dx^3}[a_0(x)T_i(x)]_{x_n} C_1 \\
 & - \frac{d^3}{dx^3}[a_0(x)T_i(x)]_{x_0} A_1
 \end{aligned}$$

for  $i = -1, 0, \dots, n+1$ . (22)

and

$$\alpha = [\alpha_{-1} \alpha_0 \dots \alpha_{n+1}]^T.$$

**5. Solution procedure to find the nodal parameters**

A typical integral element in the matrix **A** is

$$\sum_{m=0}^{n-1} I_m$$

where  $I_m = \int_{x_m}^{x_{m+1}} v_i(x)r_j(x)Z(x)dx$  and  $r_j(x)$  are the quintic B-spline basis functions or their derivatives.  $v_i(x)$  are the septic B-spline weight functions or their derivatives. It may be noted that  $I_m = 0$  if  $(x_{i-4}, x_{i+4}) \cap (x_{j-3}, x_{j+3}) \cap (x_m, x_{m+1}) = \emptyset$ .

To evaluate each  $I_m$ , we employed 7-point Gauss-Legendre quadrature formula. Thus the stiffness matrix **A** is a thirteen diagonal band matrix. The nodal parameter vector  $\alpha$  has been obtained from the system  $\mathbf{A}\alpha = \mathbf{B}$  using the band matrix solution package. We have used the FORTRAN-90 program to solve the boundary value problems (1) - (2) by the proposed method.

**6. Numerical results**

To demonstrate the applicability of the proposed method for solving the fifth order boundary value problems of the type (1) and (2), we considered two linear and two nonlinear boundary value problems. The obtained numerical results for each problem are presented in tabular forms and compared with the exact solutions available in the literature.

**Example 1:** Consider the linear boundary value problem

$$\begin{aligned}
 y^{(5)} + xy &= (1-x)\cos x - 5\sin x + x\sin x \\
 -x^2 \sin x, & 0 < x < 1
 \end{aligned}$$

(23)

subject to  
 $y(0) = y(1) = 0, y'(0) = 1, y'(1) = -\sin 1, y''(0) = -2$ .

The exact solution for the above problem is  $y = (1-x)\sin x$ . The proposed method is tested on this problem where the domain  $[0, 1]$  is divided into 10 equal subintervals. The obtained numerical results for this problem are given in Table 1. The maximum absolute error obtained by the proposed method is  $2.366304 \times 10^{-5}$ .

**Table 1:** Numerical results for Example 1

x	Absolute error by the proposed method
0.1	2.086163E-06
0.2	2.592802E-06
0.3	1.095235E-05
0.4	8.106232E-06
0.5	2.366304E-05
0.6	1.525879E-05
0.7	2.045929E-05
0.8	3.412366E-06
0.9	4.082918E-06

**Example 2:** Consider the linear boundary value problem

$$\begin{aligned}
 y^{(5)} + y^{(4)} + e^{-2x} y &= e^{-x} [-4e^{2x}(-3+x)\cos x \\
 -(1-x + 4e^{2x}(5+2x))\sin x], & 0 \leq x \leq 1
 \end{aligned}$$

(24)

subject to

$$y(0) = y(1) = 0, y'(0) = -1, y'(1) = e\sin 1, y''(0) = 0.$$

The exact solution for the above problem is  $y = e^{-x}(x-1)\sin x$ .

The proposed method is tested on this problem where the domain  $[0, 1]$  is divided into 10 equal subintervals. The obtained numerical results for this problem are given in Table 2. The maximum absolute error obtained by the proposed method is  $6.532669 \times 10^{-5}$ .

**Table 2:** Numerical results for Example 2

x	Absolute error by the proposed method
0.1	1.128763E-05
0.2	1.668930E-06
0.3	4.512072E-05
0.4	2.625585E-05
0.5	6.532669E-05
0.6	2.828240E-05
0.7	4.631281E-05
0.8	4.649162E-06
0.9	8.195639E-06

**Example 3:** Consider the nonlinear boundary value problem

$$y^{(5)} + 24e^{-5y} = \frac{48}{(1+x)^5}, \quad 0 \leq x \leq 1$$

(25)

subject to

$$y(0) = 0, y(1) = \ln 2, y'(0) = 1, y'(1) = 0.5, y''(0) = -1.$$

The exact solution for the above problem is  $y = \ln(1+x)$ .

The nonlinear boundary value problem (25) is converted into a sequence of linear boundary value problems generated by quasilinearization technique [14] as

$$y_{(n+1)}^{(5)} - 120e^{-5y_{(n)}} y_{(n+1)} = \frac{48}{(1+x)^5} - 120y_{(n)} e^{-5y_{(n)}}$$

(26)

$$-24e^{-5y_{(n)}}, n = 0, 1, 2, \dots$$

subject to

$$y_{(n+1)}(0) = 0, y_{(n+1)}(1) = \ln 2, y'_{(n+1)}(0) = 1, \\ y'_{(n+1)}(1) = 0.5, y''_{(n+1)}(0) = -1.$$

Here  $y_{(n+1)}$  is the  $(n+1)^{th}$  approximation for  $y(x)$ . The domain  $[0, 1]$  is divided into 10 equal subintervals and the proposed method is applied to the sequence of linear problems (26). The obtained numerical results for this problem are presented in Table 3. The maximum absolute error obtained by the proposed method is  $3.463030 \times 10^{-5}$ .

**Table 3:** Numerical results for Example 3

$x$	Absolute error by the proposed method
0.1	1.557171E-06
0.2	5.826354E-06
0.3	6.437302E-06
0.4	2.211332E-05
0.5	2.393126E-05
0.6	3.463030E-05
0.7	2.324581E-05
0.8	2.026558E-05
0.9	3.814697E-06

**Example 5:** Consider the nonlinear boundary value problem

$$y^{(5)} + [y']^2 e^{4y} - 4y^2 e^{y''} + e^{2x} [y''']^2 \\ = 32e^{-2x}, \quad 0 \leq x \leq 1 \quad (27)$$

subject to

$$y(0) = 1, y(1) = e^{-2}, y'(0) = -2, y'(1) = -2e^{-2}, y''(0) = 4.$$

The exact solution for the above problem is  $y = e^{-2x}$ .

The nonlinear boundary value problem (27) is converted into a sequence of linear boundary value problems generated by quasilinearization technique [14] as

$$y_{(n+1)}^{(5)} + 2e^{2x} y_{(n)}''' y_{(n+1)}''' - 4y_{(n)}^2 e^{y_{(n)}''} y_{(n+1)}'' \\ + 2y_{(n)}' e^{4y_{(n)}} y_{(n+1)}' + [4y_{(n)}'^2 e^{4y_{(n)}} - 8y_{(n)} e^{y_{(n)}''}] y_{(n+1)} \\ = 32e^{-2x} + e^{2x} y_{(n)}'''^2 - 4y_{(n)}^2 e^{y_{(n)}''} (1 + y_{(n)}'') \\ + y_{(n)}'^2 e^{4y_{(n)}} (1 + 4y_{(n)}), \quad n = 0, 1, 2, \dots \quad (28)$$

subject to

$$y_{(n+1)}(0) = 0, y'_{(n+1)}(0) = -2, y_{(n+1)}(1) = e^{-2},$$

$$y'_{(n+1)} = -2e^{-2}, y''_{(n+1)}(0) = 4.$$

Here  $y_{(n+1)}$  is the  $(n+1)^{th}$  approximation for  $y(x)$ . The domain  $[0, 1]$  is divided into 10 equal subintervals and the proposed method is applied to the sequence of linear problems (28). The obtained numerical results for this problem are presented in Table 4. The maximum absolute error obtained by the proposed method is  $8.052588 \times 10^{-5}$ .

**Table 4:** Numerical results for Example 4

$x$	Absolute error by the proposed method
0.1	3.993511E-06
0.2	2.473593E-05
0.3	4.440546E-05
0.4	6.681681E-05
0.5	8.052588E-05
0.6	7.990003E-05
0.7	6.411970E-05
0.8	4.020333E-05
0.9	1.385808E-05

## 7. Conclusions

In this paper, we have employed a Petrov-Galerkin method with quintic B-splines as basisfunctions and septic B-splines as weight functions to solve fifth order boundary value problems with special case of boundary conditions. The quintic B-spline basis set has been redefined into a new set of basis functions which vanish on the boundary where the Dirichlet boundary conditions are prescribed. The septic B-splines are redefined into a new set of weight functions which in number match the number of redefined set of basis functions. The proposed method has been tested on two linear and two nonlinear fifth order boundary value problems. The numerical results obtained by the proposed method are in good agreement with the exact solutions available in the literature. The strength of the proposed method lies in its easy applicability, accurate and efficient to solve fourth order boundary value problems.

## References

- [1] A.R.Davies, A.Karageorghis and T.N.Phillips, "Spectral Galerkin methods for the primary two-point boundary value problem in modelling viscoelastic flows", International Journal for Numerical Methods in Engineering, XVI, pp. 647-662, 1988.
- [2] H.N.Caglar, S.H.Caglar and E.H.Twizell, "The numerical solution of fifth order boundary value problems with sixth degree B-spline functions", Applied Mathematics Letters, XII, pp. 25-30, 1999.
- [3] R.P. Agarwal, Boundary Value Problems for Higher Order Differential Equations, World Scientific, Singapore, 1986.
- [4] Abdul-Majid Wazwaz, "The numerical solution of fifth order boundary value problems by the Decomposition method", Journal of Computational and Applied Mathematics, CXIVL, pp. 259-270, 2001.
- [5] Shahidi S. Siddiqi, Ghazala Akram and Arfa Elahi, "Quartic spline solution of linear fifth order boundary value problems, Applied Mathematics and Computation", CIVC, pp. 214-220, 2008.
- [6] J.Rashidinia, R.Jalilian and K.Farajeyam, "Spline approximate solution of fifth order boundary value problem, Applied Mathematics and Computation", CVIIC, pp. 107-112, 2007.

- [7] Muhammad Aslam Noor and Syed Tauseef Mohyud-Din, "An efficient algorithm for solving fifth-order boundary value problems", *Mathematical and Computer Modelling*, VL, pp. 954-964, 2007.
- [8] Hikmet Caglar and Nazan Caglar, "Solution of fifth-order boundary value problems by using Local polynomial regression", *Applied Mathematics and Computation*, CXIVC, pp. 952-956, 2007.
- [9] Mohamed El-Gamel, "Sinc and the numerical solution of fifth order boundary value problems", *Applied Mathematics and Computation*, CXIIIC, pp. 1417-1433, 2007.
- [10] Muhammed I. Syam and Basem S.Ahili, "Numerical solution of singularly perturbed fifth order two point boundary value problem", *Applied mathematics and Computation*, CXXXC, pp. 1085-1094, 2005.
- [11] Zhao-Chunwu, "Approximate analytical solutions of fifth order boundary value problems by the Variational iteration method", *Computer and Mathematics with Applications*, LXVIII, pp. 2514-2517, 2009.
- [12] A.Lamni, H.Mraoui, D.Sbibih and A.Tijini, "Sextic spline solution of fifth order boundary value problems", *Mathematics and Computers Simulation*, LXXVII, pp. 237-246, 2008.
- [13] K.N.S.Kasi Viswanadham and Sreenivasulu Ballem, "Numerical solution of fifth-order boundary value problems by Galerkin method with quartic B-splines", *International Journal of Computers Applications*, LXXVII, pp. 7-12, 2013.
- [14] R.E.Bellman and R.E. Kalaba, *Quasilinearization and Nonlinear Boundary Value Problems*, American Elsevier, New York, 1965.
- [15] L.Bers, F.John and M.Schechter, *Partial Differential Equations*, John Wiley Inter science, New York, 1964.
- [16] J.L.Lions and E.Magenes, *Non-Homogeneous Boundary Value Problem and Applications*. Springer-Verlag, Berlin, 1972.
- [17] A.R.Mitchel and R.wait, *The Finite Element Method in Partial Differential Equations*, John Wiley and Sons, London, 1997.
- [18] P.M. Prenter, *Splines and Variational Methods*, John-Wiley and Sons, New York, 1989.
- [19] Carl de-Boor, *A Pratical Guide to Splines*, Springer-Verlag, 1978.
- [20] I.J. Schoenberg, "On Spline Functions", MRC Report 625, University of Wisconsin, 1966.

### Author Profile



**S M Reddy** received the M.Sc. and Ph.D. degrees in Mathematics & scientific computing and Finite element methods from National Institute of Technology Warangal in 2007 and 2016, respectively. During 2007-2012, worked as a Assistant professor in Sreenidhi institute of science and techonology, India. During 2013-2016, worked as a research scholar in NIT Warangal, India. Now working as a Assistant professor in Sreenidhi institute of science and techonology, India.